



MERZ CONNECT[™] Support Line Insurance Verification Request Form

Please Fax the Completed Form to: 1-866-471-3005

You may contact the MERZ CONNECT[™] Support Line by calling **1-844-4MYMERZ (1-844-469-6379)**

Monday through Friday between the hours of 8am-8pm ET

PROVIDER INFORMATION		
Provider Name:	Specialty:	
Provider Email Address:		
Facility Name:	Office Contact Name:	
Street Address:		
City:	State:	Zip:
Phone # (include ext):	Secure Fax #:	May we contact your patient? Yes <input type="radio"/> No <input type="radio"/>
Place of service: (check all to be verified)		
<input type="checkbox"/> Physician Office (11)	<input type="checkbox"/> Hospital Outpatient Department (22)	
<input type="checkbox"/> Hospital Inpatient (21)	<input type="checkbox"/> Skilled Nursing Facility (31)	
<input type="checkbox"/> Ambulatory Surgical Center (21)		
<input type="checkbox"/> Other	If other, please specify: _____	
How do you intend to supply the medication?		
<input type="radio"/> Specialty Pharmacy	Name: _____	<input type="radio"/> Buy and Bill
Place of Service Name and Address (if different than Facility listed above):		
National Provider Identifier #:		State License #:
Medicare Provider #:	Medicaid Provider #:	Blue Cross Provider #:
Tax ID#:	DEA #:	Provider Identification Number (PIN):
Would you like us to determine financial assistance options for your patient and initiate the appropriate steps?	<input type="radio"/> Yes <input type="radio"/> No	
If the patient's insurer requires a prior authorization would you like assistance with pursuing the prior authorization?	<input type="radio"/> Yes <input type="radio"/> No	
<p>The MERZ CONNECT Support Line (Support Line) has been developed by Merz. The Support Line is operated for Merz by Covance Market Access Services (Covance) to help ensure every patient needing XEOMIN[®] therapy has access to reimbursement assistance. A representative of the Support Line may contact you with information relating to reimbursement assistance. By checking "Yes" below, you are certifying that the described therapy is medically necessary and you have received the necessary authorization to release the referenced medical and/or other patient information relating to XEOMIN therapy to Covance for the purpose of seeking reimbursement for XEOMIN therapy and/or assisting in initiating or continuing XEOMIN therapy, seeking copayment assistance for financially needy patients, and seeking assistance for uninsured and underinsured patients. Aggregate data regarding research requests for you and others may be shared with Merz.</p> <p>Provider Consent: <input type="radio"/> Yes <input type="radio"/> No</p>		

Please see Important Safety Information, including BOXED WARNING, on next page. Visit XEOMIN.com for Full Prescribing Information.

PATIENT INFORMATION		
<small>(Please attach an enlarged copy of the front and back of the patient's insurance card and/or other insurance information along with this form)</small>		
Patient First Name:	Patient Last Name:	MI
Patient Social Security #:	Date of Birth (mm/dd/yyyy):	Gender: F <input type="radio"/> M <input type="radio"/>
Street Address:		
City:	State:	Zip:
Daytime Phone # :		Evening Phone #:
Primary Insurance Name:		Phone #:
Subscriber Name:	Subscriber ID#:	Group ID#:
Subscriber Date of Birth (mm/dd/yyyy):	Subscriber Social Security#:	Employer Name:
Secondary Insurance Name:		Phone #:
Subscriber Name:	Subscriber ID#:	Group ID#:
Subscriber Relationship to Patient	Subscriber Social Security#:	Employer Name:
Subscriber Date of Birth (mm/dd/yyyy):		
TREATMENT INFORMATION		
Diagnosis Code 1 (ICD-10)		CPT Code 1
Diagnosis Code 2 (ICD-10)		CPT Code 2
<input type="checkbox"/> 95873 Electrical stimulation for guidance in conjunction with chemodeneration code <input type="checkbox"/> 95874 Needle electromyographic guidance (EMG) in conjunction with chemodeneration code		
<input type="checkbox"/> 76942 Ultrasound guidance for needle placement (eg, biopsy, aspiration, injection, localization device), imaging supervision, and interpretation		
Injection Sites _____		
# of Units Anticipated _____		
Estimated date of injection: _____		

IMPORTANT NOTICE: Using the MERZ CONNECT Support Line is not a guarantee of insurance benefits or payment. All benefits are subject to the insured's plan and verifications performed by Merz's third-party administrator (Covance) are not a substitute for written guidance from the plan. Under no circumstances shall the Support Line or Merz Pharmaceuticals, LLC or any Merz affiliate be held responsible or liable for payment of any claims, benefits, or costs. The practitioner must determine, based on independent medical judgment, to use XEOMIN for the specified treatment or his or her patient(s) and must supply the appropriate diagnostic codes to the Support Line for the treatment provided. It is the practitioner's sole responsibility to determine appropriate codes, charges, and modifiers, and to submit bills for services and products consistent with what was rendered as well as the patient's insurance requirements. Third-party payer coverage, reimbursement policies, and coding requirements may change over time. Providers are encouraged to contact third-party payers independently to verify specific information on their policies.

This fax was sent by Covance Market Access Services Inc, 9801 Washington Blvd., Ninth Floor, Gaithersburg, MD 20878, on behalf of Merz Pharmaceuticals, LLC, 6601 Six Forks Road, Suite 430, Raleigh, NC 27615. If you no longer wish to receive communication from Merz Pharmaceuticals, LLC, please email merzneurosciences@merz.com with subject line "Unsubscribe Merz Pharmaceuticals." If you no longer wish to receive communication from Covance, please email merzneurosciences@merz.com with subject line "Unsubscribe Covance."

We respect your privacy. To view Merz Pharmaceuticals, LLC privacy policy, visit www.merzusa.com/privacypolicy. US-XET-200

IMPORTANT SAFETY INFORMATION

INDICATIONS

XEOMIN® (incobotulinumtoxinA) for injection is indicated for the treatment of:

- Chronic sialorrhea in patients 2 years of age and older
- Upper limb spasticity in adults
- Upper limb spasticity in pediatric patients 2 years of age and older, excluding spasticity caused by cerebral palsy
- Cervical dystonia in adults
- Blepharospasm in adults

WARNING: DISTANT SPREAD OF TOXIN EFFECT
See full prescribing information for complete BOXED WARNING.

The effects of XEOMIN and all botulinum toxin products may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults, particularly in those patients who have underlying conditions that would predispose them to these symptoms.

CONTRAINDICATIONS

- Known hypersensitivity to any botulinum toxin product or to any of the components in the formulation.
- Infection at the proposed injection site(s) because it could lead to severe local or disseminated infection.

WARNINGS AND PRECAUTIONS

- The potency units of XEOMIN are specific to the preparation and assay method used and are not interchangeable with other preparations of botulinum toxin products. Therefore, Units of biological activity of XEOMIN cannot be compared to or converted into Units of any other botulinum toxin products.
- Serious hypersensitivity reactions have been reported with botulinum toxin products (anaphylaxis, serum sickness, urticaria, soft tissue edema, and dyspnea). If serious and/or immediate hypersensitivity reactions occur, discontinue further injection of XEOMIN and institute appropriate medical therapy immediately. The use of XEOMIN in patients with a known hypersensitivity to any botulinum neurotoxin or to any of the excipients (human albumin, sucrose), could lead to a life-threatening allergic reaction.
- Treatment with XEOMIN and other botulinum toxin products can result in swallowing or breathing difficulties. Patients with pre-existing swallowing or breathing difficulties may be more susceptible to these complications. When distant effects occur, additional respiratory muscles may be involved. Patients may require immediate medical attention should they develop problems with swallowing, speech, or respiratory disorders. Dysphagia may persist

for several months, which may require use of a feeding tube. Aspiration may result from severe dysphagia [See **BOXED WARNING**].

- Individuals with peripheral motor neuropathic diseases, amyotrophic lateral sclerosis, or neuromuscular junctional disorders (e.g., myasthenia gravis or Lambert-Eaton syndrome) may be at increased risk for severe dysphagia and respiratory compromise from typical doses of XEOMIN.
- **Cervical Dystonia:** Treatment with botulinum toxins may weaken neck muscles that serve as accessory muscles of ventilation. This may result in critical loss of breathing capacity in patients with respiratory disorders who may have become dependent upon these accessory muscles. There have been post-marketing reports of serious breathing difficulties, including respiratory failure, in patients with cervical dystonia treated with botulinum toxin products. Patients with smaller neck muscle mass and patients who require bilateral injections into the sternocleidomastoid muscles are at greater risk of dysphagia. Limiting the dose injected into the sternocleidomastoid muscle may decrease the occurrence of dysphagia.
- **Blepharospasm:** Injection of XEOMIN into the orbicularis oculi muscle may lead to reduced blinking and corneal exposure with possible ulceration or perforation. To decrease the risk for ectropion, XEOMIN should not be injected into the medial lower eyelid area.
- XEOMIN contains human serum albumin. Based on effective donor screening and product manufacturing processes, it carries an extremely remote risk for transmission of viral diseases and variant Creutzfeldt-Jakob disease (vCJD). There is a theoretical risk for transmission of Creutzfeldt-Jakob disease (CJD), but if that risk actually exists, the risk of transmission would also be considered extremely remote. No cases of transmission of viral diseases, CJD, or vCJD have ever been reported for albumin.

ADVERSE REACTIONS

The most commonly observed adverse reactions at rates specified below and greater than placebo are:

- **Chronic Sialorrhea:**
 - **in adults** ($\geq 4\%$ of patients): tooth extraction, dry mouth, diarrhea, and hypertension.
 - **in pediatric patients** ($\geq 1\%$ of patients): bronchitis, headache, and nausea/vomiting.
- **Upper Limb Spasticity:**
 - **in adults** ($\geq 2\%$ of patients): seizure, nasopharyngitis, dry mouth, and upper respiratory tract infection.
 - **in pediatric patients** ($\geq 3\%$ of patients): nasopharyngitis and bronchitis.
- **Cervical Dystonia in adults** ($\geq 5\%$ of patients): dysphagia, neck pain, muscle weakness, injection site pain, and musculoskeletal pain.
- **Blepharospasm in adults** ($\geq 10\%$ of patients): eyelid ptosis, dry eye, visual impairment, and dry mouth.

DRUG INTERACTIONS

Co-administration of XEOMIN and aminoglycoside or other agents interfering with neuromuscular transmission, (e.g., muscle relaxants), should only be performed with caution as these agents may potentiate the effect of the toxin.

Use of anticholinergic drugs after administration of XEOMIN may potentiate systemic anticholinergic effects.

The effect of administering different botulinum toxin products at the same time or within several months of each other is unknown. Excessive neuromuscular weakness may be exacerbated by administration of another botulinum toxin prior to the resolution of the effects of a previously administered botulinum toxin.

USE IN PREGNANCY

There are no adequate data on the developmental risk associated with the use of XEOMIN in pregnant women. XEOMIN should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

PEDIATRIC USE

Safety and effectiveness of XEOMIN in patients less than 18 years of age have not been established for lower limb spasticity, cervical dystonia, or blepharospasm.

Safety and effectiveness have been established in pediatric patients 2 to 17 years of age in patients with chronic sialorrhea and upper limb spasticity. A pediatric assessment for XEOMIN in upper limb spasticity demonstrates that XEOMIN is safe and effective in another pediatric population. However, XEOMIN is not approved for such patient population due to marketing exclusivity for another botulinum toxin.

Copyright © 2020 Merz Pharmaceuticals, LLC. All rights reserved. MERZ, the MERZ logo, and XEOMIN are registered trademarks of Merz Pharma GmbH & Co. KGaA.